

EXTRAORDINARY SAVINGS
FOR YOUR EXTRAORDINARY PET

RECEIVE \$15 REBATE FOR 2 DOSES
OR \$35 REBATE FOR 4 DOSES

TERMS AND CONDITIONS:

Offer may only be redeemed by the pet owner. Must be 18 years or older to redeem. By registering for the program you consent to opt-in to the program. You may unsubscribe at any time. To receive a rebate, all fields must be completed. A detailed invoice with purchase price must be mailed in with a copy of the rebate form. Rebate submission, including invoice, must be received by January 31, 2018. Invoice must reflect a Two (2) dose or Four (4) dose purchase of Bravecto (Fluralaner)* and/or Bravecto (Fluralaner topical solution)*. Offer valid for purchases made during the program period, between January 1, 2017, and December 31, 2017. Rebate check amount will not exceed the amount of the qualifying purchase. Purchases from online pharmacies or other retail outlets not associated with your veterinary clinic are not valid. Merck Animal Health reserves the right to decline submissions for purchases from unauthorized distributors. Void where prohibited by law. Please allow approximately 6 to 8 weeks for rebate check delivery. Unless expressly prohibited by law, payee authorizes reasonable dormancy fees deducted if check not cashed within 180 days. For rebate inquiries call 855-496-0146. May not be combined with any other offers. Offer available on purchases from clinics in the fifty (50) United States or the District of Columbia only. MERCK ANIMAL HEALTH RESERVES THE RIGHT TO CHANGE OR WITHDRAW THE PROGRAM AT ANY TIME.

PLEASE SEE FULL PRESCRIBING INFORMATION WITHIN PRODUCT PACKAGING.
TO LEARN MORE ABOUT BRAVECTO, VISIT BRAVECTO.COM.

*Bravecto kills fleas and prevents flea infestations. **Bravecto Chew for Dogs** kills ticks (black-legged tick, American dog tick, and brown dog tick) for 12 weeks and also kills lone star ticks for 8 weeks. **Bravecto Topical for Cats** kills ticks (black-legged tick) for 12 weeks and American dog ticks for 8 weeks.

Copyright © 2017 Intervet Inc., d/b/a Merck Animal Health, a subsidiary of Merck & Co., Inc. All rights reserved. US/BRV/1016/0101

BRAVECTO[®]
(FLURALANER)
BRAVECTO[®]
(FLURALANER)
TOPICAL SOLUTION

12 **TWELVE-WEEK**
PROTECTION

KEEP YOUR PET FREE OF
FLEAS AND TICKS.



How it works:

- 1 **Purchase** your Bravecto* from your veterinarian.
- 2 **Complete** the information listed below.
- 3 **Enclose** the original sales invoice with purchase price(s) circled, and the original rebate form completed from your veterinarian. **Important:** Keep copies of all materials submitted for your own records.
- 4 **Mail** your completed rebate form and requested information above to:
Merck Rebate Center
Offer Code: BRAV201701
P.O. Box 426027
Del Rio, TX 78842-6027

Name[†]: _____
Street Address[†]: _____
City[†]: _____
State[†]: _____ Zip[†]: _____
Phone Number[†]: _____
Email[†]: _____
Clinic[†]: _____
Pet Name[†]: _____
Species[†]: _____ DOB[†]: _____



***Must be completed to qualify for rebate.**

Check here if you would like to receive more information on Merck Animal Health products.

*Bravecto kills fleas and prevents flea infestations. **Bravecto Chew for Dogs** kills ticks (black-legged tick, American dog tick, and brown dog tick) for 12 weeks and also kills lone star ticks for 8 weeks. **Bravecto Topical for Cats** kills ticks (black-legged tick) for 12 weeks and American dog ticks for 8 weeks.

IMPORTANT SAFETY INFORMATION: Bravecto has not been shown to be effective for 12-weeks' duration in puppies or kittens less than 6 months of age. **Bravecto Chew for Dogs:** The most common adverse reactions recorded in clinical trials were vomiting, decreased appetite, diarrhea, lethargy, polydipsia, and flatulence. Bravecto is not effective against lone star ticks beyond 8 weeks of dosing. **Bravecto Topical for Cats:** The most common adverse reactions recorded in clinical trials were vomiting, itching, diarrhea, hair loss, decreased appetite, lethargy, and scabs/ulcerated lesions. Bravecto is not effective against American dog ticks beyond 8 weeks of dosing. For topical use only. Avoid oral ingestion. The safety of Bravecto has not been established in breeding, pregnant and lactating cats. Use with caution in cats with a history of neurologic abnormalities. Neurologic abnormalities have been reported in cats receiving Bravecto, even in cats without a history of neurologic abnormalities.

 **MERCK**
Animal Health